

DISTRIBUTION OF PESTICIDE (ORGANOCHLORINE) IN TERENGGANU
RIVER ESTUARINE SEDIMENTS

NORBADARIAH BINTI REDZUAN

FACULTY OF MARITIME STUDIES AND MARINE SCIENCE
UNIVERSITI MALAYSIA TERENGGANU
2008

Perpustakaan Sultanah Nur Zahirah (UMT)
Universiti Malaysia Terengganu

1100061803

LP 26 FMSM 2 2008



1100061803

Distribution of pesticide (organochlorine) in Terengganu river estuarine sediments / Norbadariah Redzuan.



PERPUSTAKAAN SULTAH NUR ZAHRAH
UNIVERSITI MALAYSIA TERENGGANU (UMT)
21030 KUALA TERENGGANU

11000618P3

1100061803

Lihat sebelah

HAK KILIK
PERPUSTAKAAN SULTANAH NUR ZAHIRAH UTM

**DISTRIBUTION OF PESTICIDE (ORGANOCHLORINE) IN TERENGGANU
RIVER ESTUARINE SEDIMENTS**

By

Norbadariah Binti Redzuan

Research report submitted in partial fulfillment of
the requirements for the degree of
Bachelor of Science (Marine Science)

Department of Marine Science
Faculty of Maritime Studies and Marine Science
UNIVERSITI MALAYSIA TERENGGANU

2008

This project should be cited as:

Norbadariah, R. 2008. Distribution of Pesticide (Organochlorine) in Terengganu River Estuarine Sediments. Undergraduate thesis, bachelor of Science (Marine Science), Faculty of Maritime Studies and Marine Science, University Malaysia Terengganu, Terengganu. 75p.

No part of this project report may be reproduced by any mechanical, photographic, or electronin process, or in the form of phonographic recording, or may it be stored in a retrieval system, transmitted or otherwise copied for public or private use, without written permission from the author and the supervisor of the project.

1100061803



**JABATAN SAINS MARIN
FAKULTI PENGAJIAN MARITIM DAN SAINS MARIN
UNIVERSITI MALAYSIA TERENGGANU**

**PENGAKUAN DAN PENGESAHAN SARANAN
PROJEK PENYELIDIKAN I DAN II**

Adalah ini diakui dan disahkan bahawa laporan penyelidikan bertajuk: Distribution of Pesticide (Organochlorine) in Terengganu River Estuarine Sediments oleh Norbadariah binti Redzuan, No. Matrik UK 12195 telah diperiksa dan semua pembetulan yang disarankan telah dilakukan. Laporan ini dikemukakan kepada Jabatan Sains Marin sebagai memenuhi sebahagian daripada keperluan memperolehi Ijazah Sarjana Muda Sains (Sains Samudera), Fakulti Pengajian Maritim dan Sains Marin, Universiti Malaysia Terengganu.

Disahkan oleh:

Penyelia Utama

Nama: Prof Madya Dr. Mohamed Kamil b. Abd. Rashid

Cop Rasmi:

PROF. MADYA DR. MOHAMED KAMIL ABDUL RASHID
Timbalan Dekan (Siswazah & Penyelidikan)
Fakulti Pengajian Maritim dan Sains Marin
Universiti Malaysia Terengganu (UMT)
21030 Kuala Terengganu.

Tarikh: **4.8.2008**

Ketua jabatan sains Marin

Nama: Dr. Razak bin Zakariya

Cop Rasmi:

DR. RAZAK ZAKARIYA
Ketua Jabatan Sains Marin
Fakulti Pengajian Maritim dan Sains Marin
Universiti Malaysia Terengganu
(UMT)

Tarikh: **14/5/08**

ACKNOWLEDGEMENT

In this opportunity, I would like to thank to supervisor Prof Madya Dr. Mohamed Kamil bin Abd. Rashid. It was very valuable experience during being under his supervise. Experience in doing the project independently and searching for the additional component to finish up this project with the minimum help.

More over, I also would like to say the same thing to other candidate who was done the familiar project with me. After doing several discussions, we were working together in the lab and always help each other. I hope their projects also succeed.

Last but not least, thank to mum and dad who were very supportive in giving me advice. They were understood so much within the time I was running the project. Lab assistance and science officer also help me out along the journey on sampling and sample analysis. Thank you all.

NORBADARIAH BINTI REDZUAN

UK 12195

BACHELOR OF SCIENCE (MARINE SCIENCE) (2005-2008)

TABLE OF CONTENT

CONTENT	PAGE
Acknowledgement	iii
Table of Content	iv
List of Tables	vii
List of Figures	viii
List of Abbreviation	ix
List of Symbols	x
List of Appendices	xi
Abstract	xii
Abstrak	xiii

CHAPTER 1

1.0 INTRODUCTION	1
-------------------------	---

CHAPTER 2

2.0 LITERATURE REVIEW	
2.1 Introduction to pesticide	5
2.2 Risks of Organochlorine usage as Pesticide	6
2.3 Importance of estuary	7
2.4 Pesticide usage and the route of exposure	8
2.5 Pesticide	10
2.5.1 Alpha BHC	10
2.5.2 Heptachlor and Heptachlor Epoxide	12
2.5.3 Aldrin and Dieldrin	14

2.5.4	Delta-BHC	17
2.5.5	Beta – BHC	18
2.5.6	Gamma-BHC	20
2.5.7	Endosulfan I (alpha)	22
2.5.8	4,4-DDE	24
2.5.9	4,4-DDD	25
2.5.10	4,4-DDT	26
2.5.11	Endosulfan sulfate	28
2.5.12	Endosulfan II (beta)	30
2.5.13	Endrin	31
2.5.14	Endrin aldehyde	33

CHAPTER 3

3.0 METHODOLOGY

3.1	Study area	34
3.2	Lab preparation	36
3.3	Sediment sample collection	37
3.4	Physical parameter measurement on sampling site	38
3.5	Sample analysis	39
3.5.1	Freeze Dry	39
3.5.2	Lipid Extraction	40
3.5.3	Sample Concentration	41
3.5.4	Acid Dilution	42
3.5.5	Sulphur Excretion	43
3.5.6	Silica Activation	44
3.5.7	Alimina Activation	44
3.5.8	Silica /Alumina Column	45

3.5.9	Organochlorine Analysis using GC-ECD	46
3.6	Total Organic Carbon Analysis	50

CHAPTER 4

4.0 RESULT

4.1	Result of OCP concentration	50
4.2	Lipid percentage and abundance of pesticide	52
4.3	Pesticide response on physical parameter	53
4.4	Result Total Organic In Sediment Sample	56
4.5	Recovery Test	57

CHAPTER 5

DISCUSSION	58
-------------------	----

CHAPTER 6

CONCLUSION	62
-------------------	----

REFERENCES

APPENDICES

VITAE CURICULUM

LIST OF TABLE

TABLE	PAGE
3. 1 : Coordinate of sampling station.	35
4.2 : Result of glucose percentage in TOC analysis.	56
4. 3 : Percentage of recovery test according the OCP species.	57

LIST OF FIGURES

FIGURE	PAGE
3. 1 : Map of Terengganu River Estuary and location for sampling stations.	35
4. 1 : Relation between concentration of species gamma-BHC and in surface DO.	52
4. 2 : Linear relation between concentration of alpha-BHC against water depth.	54
4. 3 : Linear relation between concentration of delta-BHC against water depth.	54
4. 4 : Linear relation between concentration of heptachlor epoxide against water depth.	55
4. 5 : Linear relation between concentration of DDE against water depth.	55
4.6 : Linear relation between concentration of DDE against water depth.	56

LIST OF ABBREVIATION

BHC	: Benzene hexachloride
DCM	: Dichloromethane
DDD	: Dichloro-Diphenyl-Dichloroethane
DDE	: Dichloro-Diphenyldichloro-Ethylene
DDT	: Dichloro-Diphenyl-Trichloroethane
DO	: Dissolved oxygen
GC-ECD	: Gas Chromatography-electron capture detector
GCIS	: Gas Chromatography Internal Standard
GPS	: Global Positioning System
HCH	: Hexachlorocyclohexane
HCl	: Hydrochloric acid
Na ₂ SO ₄	: Sodium sulphate
ND	: Not detected
OCP	: Organochlorine pesticide
PCB	: Polychlorinated biphenyl
ppt	: Part per thousand
TCMX	: Tetra- chloro-m-xylene
TEL	: Total Extractable Lipid

LIST OF SYMBOLS

μ	: micron
μL	: micronlitre
$^{\circ}\text{C}$: degree Celsius
g	: gram
kg	: kilogram
L	: litre
mL	: mililitre
M	: mole
ng	: nanogram
sec	: second
V	: volume

LIST OF APPENDICES

APPENDICES	PAGES
APPENDIX 1: Concentration alpha-BHC in sediment sample.	67
APPENDIX 2: Two-way ANNOVA calculation on 1 st sampling concentration.	83
APPENDIX 3: Figure TOC percentage abundance in 1 st sampling.	83
APPENDIX 4: Figure on TOC percentage abundance in 2 nd sampling.	84
APPENDIX 5: Salinity effect on alpha-BHC.	84
APPENDIX 6: Increasement of OCP versus surface temperature	88
APPENDIX 7: Increasement of OCP versus bottom temperature	88
APPENDIX 8: Increasement of OCP versus surface pH.	88
APPENDIX 9: Increasement of OCP versus bottom pH.	89
APPENDIX 10: Lipid Percentage in 1 st and 2 nd sampling.	89

DISTRIBUTION OF PESTICIDE (ORGANOCHLORINE) IN TERENGGANU RIVER ESTUARINE SEDIMENTS

ABSTRACT

Organochlorine pesticide in sediment samples were detected using Gas Chromatography - Electron Capture Detector (GC-ECD). The primary aim of this study is to determine the species of organochlorine pesticide (OCP) and their concentration in Terengganu River Estuarine sediment. The levels of contamination of 16 species organochlorine pesticides such as HCHs, DDTs, aldrin, dieldrin and endrin were determined in estuarine sediments along the Terengganu River Estuary in Terengganu. The concentration of various pesticides in sediments from the mouth of estuaries along the river were in the range of 0.01–1853095 ng/g (alpha-BHC), 0.002–1669 ng/g (beta BHC), 0.0002–6048 ng/g (gamma-BHC), 0.0003–2422 ng/g (delta-BHC), 0.02–17874 ng/g (heptachlor), 0.01 – 49029 ng/g (aldrin), 0.004 – 9565 ng/g (heptachlor epoxide isomer B), 0.002 – 69855 ng/g (endosulfan alpha), 0.01 – 11679280 ng/g (dieldrin), 0.15 – 5972 ng/g (DDE), 0.28 – 8428 ng/g (endrin), 0.42 – 1899 (endosulfan beta), 0.47 – 24132 ng/g (DDD), 0.5 – 2484 ng/g (endrin aldehyde), 0.14 – 4652 ng/g (endosulfan sulfate) and 21.5 – 29006 ng/g (DDT). Among the organochlorine pesticides, dieldrin and gamma-BHC were most dominant in the estuarine sediments. The highest concentration of dieldrin was estimated at station 2 near the mouth of Terengganu River estuary (11679280 ng/g) while that of gamma-BHC at the same station (6048 ng/g). Among the cyclodiene compounds, alpha-BHC was in abundance in most of the sediments whereas DDT could be detected at the smaller river channel.

ABSTRAK

Gas Chromatography - Electron Capture Detector (GC-ECD) adalah merupakan keadaan paling ideal untuk mengesan taburan dan kepekatan OCP dalam sediment. Tujuan keadaan ini dijalankan adalah untuk mengenalpasti spesis OCP yang ada di muara Sungai Terengganu beserta kepekatananya. 16 spesis OCP telah dikesan di kawasan kajian dan didapati bahawa spesis yang telah diharamkan penggunaannya juga dijumpai seperti HCH, DDT, aldrin, dieldrin adalah 0.01–1853095 ng/g 0.002–1669 ng/g (beta BHC), 0.0002–6048 ng/g (gamma dan endrin. Kepekatan pestisid yang dijumpai adalah dalam berbagai-bagai julat. Bagi sepsis alpha-BHC -BHC), 0.0003–2422 ng/g (delta-BHC), 0.02–17874 ng/g (heptachlor), 0.01 – 49029 ng/g (aldrin), 0.004 – 9565 ng/g (heptachlor epoxide isomer B), 0.002 – 69855 ng/g (endosulfan alpha), 0.01 – 11679280 ng/g (dieldrin), 0.15 – 5972 ng/g (DDE), 0.28 – 8428 ng/g (endrin), 0.42 – 1899 (endosulfan beta), 0.47 – 24132 ng/g (DDD), 0.5 – 2484 ng/g (endrin aldehyde), 0.14 – 4652 ng/g (endosulfan sulfate) and 21.5 – 29006 ng/g (DDT). Gamma-BHC dan dielrin telah dijumpai sebagai sepsis yang mendominasi kawasan kajian. Kepekatan dielrin paling tinggi telah dijumpai di stesen kajian 2 dengan kepekatan 11679280 ng/g manakala gamma-BHC ialah 6048 ng/g. Bagi spesis yang terhasil melelui tindakbalas kimia, alpha-BHC adalah senang dijumpai di dalam sample manakala DDT jarang dijumpai.