THE EFFECTS OF RETENTION PERIOD, INCUBATION PERIOD AND INCUBATION METHOD ON TURTLE HATCHLING VIGOUR

PUSHPALATHA A/P M. PALANIAPPAN

FAKULTI PERIKANAN DAN SAINS SAMUDERA UNIVERSITI PERTANIAN MALAYSIA SERDANG, SELANGOR 1994

404 1100023815

LP 48 FPSS 1 1994



The effects of retention period, incubation period and incubation method on turtle hatchling vigour / Pushpalatha M.Palaniappan.



PERPUSTAKAAN KOLEJ UNIVERSITI SAINS & TEKNOLOGI MALAYSIA 21030 KUALA TERENGGANU

	1000238.	15
		-
Part of the		
		PHOTOTOLIA III Edin Mentring and Sound
distribution in		
		Y 31 - 4 1 - 1 - 1

Lihat sebelah

HAK MILIK PERPUSTAKAAN KUSTEM

THE EFFECTS OF RETENTION PERIOD, INCUBATION PERIOD AND INCUBATION METHOD ON TURTLE HATCHLING VIGOUR

Ву

PUSHPALATHA A/P M.PALANIAPPAN

This project report is submitted as a partial requirement for the degree of Bachelor Science of Fisheries.

FAKULTI PERIKANAN DAN SAINS SAMUDERA UNIVERSITI PERTANIAN MALAYSIA SERDANG, SELANGOR 1994

PREFACE

Releasing animals, releasing turtles, is pleasurable - watching how they crawl over the sand, struggle through the surf and breaking waves and then swim out beyond to the open sea. It is not so easy to define exactly why this is pleasurable. Perhaps it has something to do with the desire for a freer simpler existence, with an element of going beyond the present situations well as back to childhood.

A.D. Graham

The Gardeners of Eden

Om Sarva vigna haram Devam Sarva vigna vivarjitham Sarva siddhi pradataram Vardeham gnana Nayakam

special thanks to Jahaten Perilbanan Teranggang as well

ACKNOWLEDGEMENTS

I am deeply indebted to Lord Muruga for the strength and perseverance in being able to complete this project. This would also be impossible without the support and encouragement from my beloved parents, Mr. and Mrs. M. Palaniappan and brother Anand as well as Ms. Lee Shiau Chin.

I would like to express my gratitude for all the guidance and advice given by both my supervisors, Mr. Liew Hock Chark and Assoc. Prof. Dr. Chan Eng Heng. Much appreciated also is all the assistance given by "Big Brother" Syed Zainuddin bin Engku Chik Abdullah, Shahbudin Saad, Roslam Abd Ghani, Ms. Ivy Wong Ling Ling, Ms. Panawon Jirawong, Mazalina bt. Ali, Abdul Talib Sulaiman, the Redang Island project pioneers, my 19 macho fellow course-mates, the third year students and all the other faculty staff. I would like to give special thanks to Jabatan Perikanan Terengganu as well the Rantau Abang Turtle Sanctuary Manager and staff as for providing the turtle hatchlings and allowing me to use their facilities. Lastly, to Muhamad Saini bin Suliansa, thank you for starting me off on the journey into this astonishing world of sea turtles.

ABSTRACT

A study on turtle hatchling vigour upon emergence in relation to parameters such as retention period, incubation period and incubation method was conducted. Six different retention periods and three methods of incubation were used in this study. Incubation period was determined by the number of days the hatchlings took to emerge from their nests. Terrestrial locomotion of the hatchlings traversing the beach from their nest to the surf was used as a measure of their vigour. A total of 24 leatherback, <u>Dermochelys</u> coriacea and 180 green turtle, Chelonia mydas hatchlings were tested. The results obtained indicated that hatchlings released immediately upon emergence had maximum vigour in comparison to those that were retained. There were significant differences in hatchling vigour in relation to the various incubation periods and incubation methods that were used in the study.

ABSTRAK

Kajian mengenai kekuatan merangkak anak penyu yang baru menetas telah dijalankan. Ianya dilakukan dengan menggunakan parameter jangkamasa kurungan, jangkamasa cara pengeraman yang berbeza. Enam pengeraman dan jangkamasa kurungan dan tiga cara pengeraman yang berbeza telah dikaji. Jangkamasa pengeraman ditentukan oleh masa yang diambil untuk kemunculan keluar anak penyu dari sarangnya. Kekuatan anak penyu diukur semasa ianya merangkak dari sarang sehingga ia sampai ke gigi air. Sejumlah 24 ekor anak penyu belimbing (Dermochelys <u>coriacea</u>) dan 180 ekor anak penyu agar (<u>Chelonia mydas</u>) telah digunakan di dalam kajian ini. Keputusan menunjukkan bahawa anak penyu yang dilepaskan sebaik sahaja ia muncul dari sarang adalah yang paling kuat jika dibandingkan dengan anak penyu yang terkurung. Perbezaan yang ketara di dalam kekuatan anak penyu telah dikesan apabila jangkamasa pengeraman dan cara pengeraman yang berlainan digunakan.