

ENDOPARASITE IDENTIFICATION IN PARROTFISH,
Siganus pruriogaster, AT MARINE PARK CENTER,
REDANG ISLAND

KIEW WOON LEY

FACULTY OF SCIENCE AND TECHNOLOGY
KOLEJ UNIVERSITI SAINS DAN TEKNOLOGI
MALAYSIA
2004

1100028945			
PERPUSTAKAAN KOLEJ UNIVERSITI SAINS & TEKNOLOGI MALAYSIA (KUSTEM)			
Pengarang <i>Kiew Woon Ley.</i>	No. Panggilan <i>dn 1693</i>		
Judul <i>Endoparasite identification in ...</i>			
Tarikh	Waktu Pemulangan	Nombor Ahli	Tanda tangan
<i>3/2/04</i>	<i>4/00pm</i>	<i>UK 6578</i>	<i>[Signature]</i>
<i>26/03/07</i>	<i>2300 hrs</i>	<i>UK10987</i>	<i>[Signature]</i>

31/3/20

1100028945

LP 8 FST 1 2004



1100028945

Endoparasit identification in parrotfish, *Scarus prasiognathos*, at marine park center, Redang Island / Kiew Woon Ley.



PERPUSTAKAAN

KOLEJ UNIVERSITI SAINS & TEKNOLOGI MALAYSIA
21030 KUALA TERENGGANU

1100028945

1100028945

Lihat sebelah

HAK MILIK
PERPUSTAKAAN KUSTEM

ENDOPARASITE IDENTIFICATION IN PARROTFISH, *Scarus prasiognathos*,
AT MARINE PARK CENTRE, REDANG ISLAND

By

KIEW WOON LEY

Research Report submitted in partial fulfillment of
the requirements of the degree of
Bachelor of Science (Marine Biology)

Department of Marine Sciences
Faculty of Science and Technology
KOLEJ UNIVERSITI SAINS DAN TEKNOLOGI MALAYSIA

2004

This project report should be cited as:

Kiew, W. L. 2004. Endoparasite identification in parrotfish, *Scarus prasiognathos*, at Marine Park Centre, Redang Island. Undergraduate thesis. Bachelor of Science in Marine Biology. Faculty of Applied Science and Technology. College University of Science and Technology Malaysia. 66p.

No part of this project report may be reproduced by any mechanical, photographic, or electronic process, or in the form of phonographic recording, nor may it be stored in a retrieval system, transmitted, or otherwise copied for public or private use, without permission from the author and supervisor of this project.

1100028945



**JABATAN SAINS SAMUDERA
FAKULTI SAINS DAN TEKNOLOGI
KOLEJ UNIVERSITI SAINS DAN TEKNOLOGI
MALAYSIA**

**PENGAKUAN DAN PENGESAHAN LAPORAN
PROJEK PENYELIDIKAN I DAN II**

Adalah ini diakui dan disahkan bahawa laporan penyelidikan bertajuk:

Endoparasite Identification in Parrotfish, *Scarus prasiognathos*, at Marine Park Centre, Redang Island oleh Kiew Woon Ley, No. Matrik UK 5576 telah diperiksa dan semua pembetulan yang disaranlan telah dilakukan. Laporan ini dikemukaan kepada Jabatan Sains Samudera sebagai memenuhi sebahagian daripada keperluan memperolehi Ijazah Sarjana Muda Sains Biologi Marin, Fakulti Sains dan Teknologi, Kolej Universiti Sains dan Teknology Malaysia.

Disahkan oleh:

Penyelia Utama

Nama : Prof. Faizah Shaharom
Cop Rasmi : PROF DR. FAIZAH SHAHAROM
Timbalan Pengarah
Institut Oseanografi
Kolej Universiti Sains dan Teknologi Malaysia
21030 Kuala Terengganu
Terengganu Darul Iman

Ketua Jabatan Sains Samudera

Nama :
Cop Rasmi :
PROF. MADYA DR. KAMARUZZAMAN B. YUNUS
Ketua
Jabatan Sains Samudera
Fakulti Sains dan Teknologi
Kolej Universiti Sains dan Teknologi Malaysia
21030 Kuala Terengganu.

ACKNOWLEDGEMENTS

I am grateful to my supervisor, Prof. Faizah Shaharom who has helped, discussed and commented critically during the progress of my project. I thank to Prof. Bjørn Berland for his advises and comments during last stage of writing. Much of this study was done in Marine Park Centre, Redang Island and I am grateful to Mr. Rahim for providing me with an opportunity to stay and work under such pleasant conditions and providing me with laboratory facilities. I thank officers and staff in Marine Park Centre for practical help during my work going on. I am grateful to my colleague, Lo Chun Nghen, giving me support and help during the progress of my work. For laboratory assistance at KUSTEM, I thank Madam Kartini and Mr. Che Zan. I am indebted to Kolej Universiti Sains dan Teknologi Malaysia, University Putra Malaysia and University Malaya for permission to use materials in library.

TABLE OF CONTENT

Content	Pages
Title Page	i
Approval form	ii
Acknowledgements	iii
Tables of Contents	v
List of Table	vi
List of Figures	viii
List of Appendices	ix
Abstract	1
Chapter 1 Introduction	1
1.1 <i>Scarus prasiognathos</i> , Valenciennes, 1840	3
1.2 Objective	4
Chapter 2 Literature Review	6
2.1 Parrotfish	6
2.2 Parasitism	7
2.2.1 Importance of parasite	7
2.2.2 Endoparasite	10
2.3 Type of endoparasite commonly found	12
2.3.1 Subclass Digenea	12
2.3.2 Subclass Cestode	12
2.3.3 Phylum Nematode	13
2.3.4 Phylum Acanthocephalan	14
2.3.5 Phylum Protozoa	15
Chapter 3 Materials and Method	16
3.1 Sampling method	16
3.2 Parasite Examination, Fixation and Staining Process	17
3.2.1 Digenea	18
3.2.2 Nematode	19
3.2.3 Acanthocephala and cestode	19
3.2.4 Protozoa	20
3.3 Drawing, Measuring and Parasites Identification	20
3.4 Calculation	21

Chapter 4 Results	22
4.1 Endoparasite found in <i>Scarus prasiognathos</i>	24
4.1.1 Digenea <i>Hapladena</i> sp.	24
4.1.2 Digenea <i>Eurycreadium</i> sp.	28
4.1.3 Myxosporidia	34
Chapter 5 Discussion	36
Chapter 6 Conclusion	40
References	42
Appendices	46
Curriculum Vitae	53

LIST OF TABLES

Table	Page
1 Water quality parameter in Marine Park Center in the period May-August 2003	22
2 Total number of endoparasites found in <i>Scarus prasiognathos</i>	23
3 Prevalence and mean intensity of endoparasites found in parrotfish, <i>Scarus prasiognathos</i>	23

LIST OF FIGURES

Figure		Page
1	Digital image of parrotfish, <i>S. prasiognathos</i>	4
2	Some examples of scolex or strobila that tapeworm, cestode, having. (source: Fernando <i>et al</i> 1972)	13
3	Some example of proboscis with its hooks in acanthocephalan (source: Fernando <i>et al</i> 1972)	14
4	Myxosporean (Source: www.fishdisease.net/fd/disease_images/image.php?img_id=218)	15
5	Internal organs of fish	18
6	Drawing of Digenea <i>Hapladena</i> sp. Scale bar = 24.4 μm	26
7	Digital image of digenea <i>Hapladena</i> sp	27
8	Drawing of Digenea <i>Eurycreadium</i> sp. Scale bar = 24.4 μm	29
9	Digital image of digenea <i>Eurycreadium</i> sp	30
10	Oral sucker of Digenea, at 20x magnification	31
11	Ventral sucker of Digenea, at 50x magnification	31

12	Digestive system of Digenea, at 25x magnification. Oesophagus was fairly long (arrow in red) and the intestine content turn into brown after preserving (arrow in black).	32
13	Genital pore of Digenea, at 20x magnification 20x. Genital pore was opens median, anterior to ventral sucker (arrow pointed)..	32
14	Reproductive system of Digenea. Two testes located opposite and ovary located between testes (with red arrow). Fertilized eggs were nearly to release.	33
15	Excretory pore of Digenea with I-shaped excretory vesicle (arrow pointed).	33
16	Drawing of Myxosporidean. Scale bar = 10.0 μm	35
17	Digital image of Myxosporidean. A: polar capsule are seen clearly in Myxosporidean; B: sutural line was almost sinuous (arrow pointed).	35

LIST OF APPENDICES

Appendix		Page
1	Location of Sampling site.	46
2	Fish sampling method	47
3	Summary of method	48
4	Endoparasites found in a total of 30 parrotfish, <i>Scarus prasiognathos</i> , in Marine Park Center, Redang Island, May-August 2003.	49
5	Water quality parameters in Marine Park Center, May - August 2003	50
6	Apparatus required	51
7	Chemicals required	52

ABSTRACT

As the important of the marine fish is being concern, more studies should be carried in order to contribute in particular field. This research had identified the endoparasite of parrotfish, (*Scarus prasiognathos*) in Marine Park Center, Redang Island. Fish were collected by handling net, identified, measured and gutted for stomach content determination. Parasite were fixed in 70 % ethanol and stained in Mayer's carmalum. Glacial Acetic Acid was used as dehydrant process. Cleared overnight in clove oil and mounted in Canada balsam. Prevalence and mean intensity of infection were calculated for each parasite species. In this study, a total of 3 difference species were found, 2 digeneans and 1 myxosporidian. With the features described, trematodes were probably identified to genera *Hapladena* sp. (with prevalence of 13.33 and mean intensity of 1.50); *Eurycreadium* sp. (with prevalence of 70.0 and mean intensity of 13.38). For the unidentified Myxosporidean the prevalence was 23.33, mean intensity 1515.29.

ABSTRAK

“Identifikasi endoparasit pada ikan payan, *Scarus prasiognathos*, di Taman Laut, Pulau Redang.”

Oleh sebab ikan air masin sangat penting, lebih banyak kajian seharusnya dijalankan supaya dapat menyumbang ke atas bidang yang berkenaan. Kajian ini telah mengenalpasti parasit-parasit yang terdapat pada ikan payan (*Scarus prasiognathos*) di Taman Laut, Pulau Redang. Ikan telah dikumpul dengan menggunakan jala tangan. Proses seperti membuat ukuran, pembedahan ikan dan identifikasi parasite telah dilakukan. Parasit dikukuhkan (fix) dengan 70 % alkohol dan diwarnakan dengan ‘Mayer’s carmalum stain’. Siri GAA diguna untuk proses dehidrasi. Parasit direndam di dalam ‘clove oil’ semalam dan akhirnya sekali spesimen dilekap dengan menggunakan ‘Canada balsam’. Setiap parasite dinilai prevalent dan min intensiti. Dalam kajian ini, sejumlah tiga jenis parasite dapat dijumpai iaitu 2 jenis digenea dan 1 jenis myxosporidia. Dengan sifat-sifat yang diuraikan, trematod berkemungkinan dari genera *Hapladena* sp. (memberi prevalent sebanyak 13.33 dan min intensity sebanyak 1.5); *Eurycreadium* sp. (memberi prevalence sebanyak 70.0 dan min intensiti sebanyak 13.38). Bagi Myxosporidean yang tidak dapat dikenalpasti telah memberi prevalence sebanyak 23.33 dan min intensiti 1515.29.